

# Behavioral Adaptations and Survival Strategies in Wild Animals: A Review

Vishesh Dubey<sup>1</sup>, Dr. Prem Chand<sup>1\*</sup> and Dr. Ajay Kumar<sup>1\*</sup>

1: Department of Zoology, \*Corresponding Author

Tilak Dhari P.G. College, Jaunpur

Affiliated to VBSPU Jaunpur



<https://doi.org/10.55041/ijst.v2i9.013>

**Cite this Article:** Dubey, V., Chand, P. & Kumar, A. (2026). Behavioral Adaptations and Survival Strategies in Wild Animals: A Review. International Journal of Science, Strategic Management and Technology, 02(9), 1-9. <https://doi.org/10.55041/ijst.v2i9.013>

**License:**  This article is published under the Creative Commons Attribution 4.0 International License (CC BY 4.0), permitting use, distribution, and reproduction in any medium, provided the original author(s) and source are properly credited.

**Abstract:** Wild animals live in environments that change continuously in temperature, rainfall, food availability, predation risk, competition, and habitat structure. Their persistence depends not only on anatomical and physiological traits but also on behavior. Behavioral adaptations are actions or patterns of activity that improve an individual's ability to survive and reproduce under particular ecological conditions. This review examines major behavioral survival strategies in wild animals, including migration, hibernation and torpor, camouflage and crypsis, antipredator behavior, territoriality, communication, social cooperation, parental care, foraging strategies, and responses to environmental disturbance. Migration permits animals to track seasonally favorable resources; dormancy reduces energetic expenditure during unfavorable periods; camouflage decreases detection; and coordinated social behavior can improve defense, hunting, and care of young. Behavioral flexibility is especially important when animals encounter rapid habitat modification, climate variability, pollution, and human disturbance. At the same time, behaviors have energetic and ecological costs and are constrained by trade-offs among feeding, reproduction, safety, and competition. Understanding these mechanisms is therefore valuable for wildlife biology and conservation management. Behavioral observations can reveal habitat quality, disturbance responses, breeding success, and changes in resource availability before population declines become obvious. The review concludes that survival in wild animals is best understood as an integrated outcome of behavior, physiology, morphology, ecology, and environmental conditions.

**Keywords:** Behavioral Adaptation; Survival Strategies; Wildlife; Migration; Camouflage; Territoriality; Social Behaviour; Conservation

**1. Introduction:** In simple words, behavior is one of the most effective ways in which different animal of the same species or individuals of the different species interacts with each other and its changing environment. As like many structural trait, behavior of an individual can sometimes change very rapidly within an individual's lifetime and can therefore provide an important mechanism for coping with environmental variability. Wild animals change their behavior continuously and that affect the decisions of individuals about where to roam, when to feed, whom to approach or avoid, when and where to reproduce, and how to respond against prey or predators or competitors. These behavioral decisions influence survival and reproductive success of individuals and, consequently, population dynamics of that species. (Alcock, 2013; Krebs & Davies, 1993).

Behavioral adaptations are highly conserved in specific animals. Different animals operate together with morphology and physiology in changing environment. As shown by different animals, migration of wild animals depends on orientation in specific direction using to identify suitable site land marker, energy storage that helps in movement during migration, locomotor capacity of the animals, and the availability of suitable habitats along a migratory route. Hibernation depends on physiological changes that lower metabolic expenditure, while behavioral selection of protected shelters reduces exposure to environmental hazards. Similarly, social cooperation may depend on communication systems, recognition of relatives, dominance relationships, and mechanisms for resolving conflict. (Davies, Krebs, & West, 2012).

The behavioral adaption in changing environment plays a significant role in survival of individual. Feeding habits of individual affect prey populations and vegetation in specific area; territorial behavior directly influences the distribution of animals in their habitat; pollination and seed dispersal connect animal movement with plant reproduction and predator behavior in natural environment regulate prey communities. Consequently, behavioral ecology provides a bridge between individual-level decisions and ecosystem-level processes. (Begon et al., 2006).

**2. Concept of Behavioral Adaptation:** A behavioral adaptation in wild animals is a behavior that increases fitness under particular environmental conditions. Fitness of the individual refers broadly to the contribution makes by an individual to the next generation through survival and reproduction. Certain characters of animal's behavior are genetically influenced and shaped over generation to generation by natural selection, while certain behavioral characters animals learn from the environment, from their experience, social transmission, or flexible responses to changing conditions. In nature real populations, behavior commonly reflects an interaction between inherited tendencies and environmental experience. (Alcock, 2013; Mayr, 2004).

There are different types of behavioral adaptations represented by wild animals in various activities. Behavioral adaptations can be classified according to their major function. Resource-acquisition behaviors help animals find food and water from the environment. Antipredator or survival from the predator behaviors reduce the probability of detection from the prey, adaptation from capture by predators, or injury (Palmer MS et al., 2021; Caro TM. 1994; LaGory KE 1987; Hasson O. 1991; Jarman P. 1974; Lima SL, Dill LM. 1990). Reproductive behaviors in wild animals including insects increase mating opportunities and maximum offspring production and their survival (Michael A Fedak et al., 2009; Kovacks K; 2009; Anholt RRH et al., 2021; Alexandra Alvergne and Vedrana Höggqvist Tabor, 2018). Social behaviors in wild animals facilitate cooperation among the intraspecies or interspecies animals, information exchange through different posture or behavior, defense, and care of young. Movement strategies, including dispersal and migration as per the environmental changes, allow animals to track resources and suitable climatic conditions. (Danchin, Giraldeau, & Cézilly, 2004).

A key feature of adaptation in wild animals is trade-off. Spending more time during feeding can increase energy intake in animals but that increased time in feeding will affect the population of predator's exposure at prey population. Defending a territory by the wild animals can secure food or mates but requires energy and may cause injury in the same (J.O. Wolff and J.A. Peterson 2010; García-Ruiz I, Rubenstein DR. 2026). Producing many offspring can increase reproductive output but can reduce parental investment per offspring (Lynch RF. 2016; Santos ESA, Nakagawa S. 2012). Thus, behavior should be interpreted as a balance among competing costs and benefits rather than as a perfect response to the environment. (Krebs & Davies, 1993; Stephens & Krebs, 1986).

**3. Migration as a Survival Strategy:** Migration is a repeated movement between geographic areas, usually associated with predictable seasonal or fixed pattern of life-history conditions in different wild animals. It is widespread among birds, mammals, fishes, insects, reptiles, and other animals. Migration allows individuals to exploit or proper utilization of temporary concentrations of food in that geographical area, help to reach breeding grounds, avoid severe climatic conditions, or complete particular stages of their life cycle. (Dingle & Drake, 2007; Fryxell, Sinclair, & Caughley, 2014).

Migratory decisions in animals are influenced by length of day or night or in other words photoperiod, temperature, rainfall, food availability, water and soil conditions, social cues, and inherited seasonal programs (Helm B, Liedvogel M. 2024; Able KP, Belthoff JR. 1998; Åkesson S, Helm B. 2020; Åkesson S, et al., 2017; Appenroth D et al., 2021). Migration either affected by environmental factors (extrinsic factors) or in some animals through affected by genetic

factors (intrinsic behavior) (Archer LC et al., 2019). Many birds use multiple environmental cues and sophisticated navigation systems. Migratory fishes may respond to chemical signals, flow conditions, temperature gradients, and geomagnetic information. Large terrestrial mammals may follow traditional routes between seasonal ranges where forage and water availability change through the year. (Berthold, 2001).

Migration carries substantial costs in living organisms throughout the migratory path. Animals must utilize their energy during movement and may face different predators, harsh environmental conditions as like storms, dehydration, disease, starvation, and anthropogenic barriers. Habitat loss at end point sites can be particularly damaging because many migrants depend on a sequence of suitable locations. Conservation therefore requires protection of breeding and wintering habitats, movement corridors, and intermediate resource sites. (Wilcove & Wikelski, 2008; Fryxell et al., 2014).

**4. Hibernation, Torpor and Dormancy:** Hibernation occurs in animals during periods of low food availability or unfavorable or harsh temperature can make ordinary activity energetically expensive. Many animals respond by reducing activity and metabolic demand. Hibernation is a prolonged inactive state involving strong reductions in body temperature and metabolic rate, while torpor a state of physically or mentally inactive condition, sluggishness, or a specialized biological condition may occur for shorter periods. Some species use daily torpor, whereas others undergo seasonal dormancy lasting weeks or months. (Geiser, 2013; Ruf & Geiser, 2015).

The behavioral decision of wild animals helps to enter a dormant state and it can conserve energy when the expected benefits of continued activity are low. Animals may prepare for the dormant condition by accumulating extra fat reserves before dormancy, selecting sheltered sites, reducing exposure to predators, and altering feeding schedules. Such characters of dormancy as like in bats, rodents, some carnivores, amphibians, reptiles, and many invertebrates show forms of metabolic depression or dormancy, although exact patterns differ among taxa (Geiser, 2013). Dormancy also involves risks. Reduced responsiveness can make an animal vulnerable to disturbance, and interruptions of torpor may consume significant energy. Climate change can alter the timing of winter conditions and food availability, creating mismatches between dormancy schedules and environmental resources. (Ruf & Geiser, 2015).

**5. Camouflage, Crypsis and Mimicry:** Camouflage, or crypsis character shows by wild animals, reduces the probability that an animal will be detected by predators or prey. Camouflage is involved matching of the background colour and pattern and it helps in hiding with predators, disruptive markings that break up body outlines, remaining motionless, or selecting backgrounds that provide visual concealment. Camouflage is particularly effective when the behavior and habitat of the animal complement its body pattern. (Ruxton, Sherratt, & Speed, 2004). Some animals also show mimicry, in which their appearance resembles another organism or environmental object to save themselves from their predator. Batesian mimicry can help a harmless species to resemble a predator like species, while mullerian mimicry involves more than one species sharing warning patterns. These strategies show how predator like appearance or perception can shape animal appearance and behavior (Ruxton et al., 2004). Camouflage is not only restricted to appearance of animals. Behavioral crypsis of the animals includes freezing, hiding, moving slowly, selecting shaded microhabitats, changing their day-by-day activity times, and avoiding exposed areas with predators. Although no camouflage is universally perfect or effective because visual backgrounds, predators, light conditions, and habitat structure vary from individual to individual.

**6. Antipredator and Defensive Behavior:** Predation of an individual animals is a major selective force shaping the behavior of individual. Animals may help to reduce risk through active vigilance, alarm calls from the other animals, high speed for escape with predators, hiding from the predator, grouping, distraction, intimidation, or active defense with predators. The wild animals choose appropriate strategy depends on their body size, habitat, predator type, distance to shelter, and the presence of offspring or social partners. (Lima & Dill, 1990; Caro, 2005). Vigilance is most important factor for the wild animals living in open habitats. Animals living social life or group-living animals can define their group through monitoring among several individuals, and it potentially allowing more time for feeding while maintaining predator detection. Alarm calls against the positive attacks from the predators can warn group members and may also communicate information about the type or location of danger. Certain wild animals' species use mobbing behavior, in which several certain individuals' approach or harass a predator (Caro, 2005). Defensive behavior of the

animals may involve frighten activity, threat postures, chemical defenses, autotomy, feigning death, or rapid escape. Many animals combine several defenses in sequence: detection, freezing, concealment, flight, and finally physical defense if escape is impossible.

**7. Territorial Behaviour and Resource Defense:** Different wild animals create their own territory demarcation through different ways. Territorial behavior occurs when animals defend an area where individual group of animals are residing, against members of the same species (intraspecies) or, in some cases, other species (inter species competition). Territories may provide access to enormous amount of food for individual species, proper nesting sites, shelter against environmental changes, water, or mating opportunities. Defense against predators can include vocal displays, scent marking using different pheromones, visual signals, patrols, and direct aggression. (Krebs & Davies, 1993). Territory defined by individual animal species, help to reduce the competition by establishing predictable access to resources. However, maintaining and defending a territory requires time and energy. Animals may therefore adjust territory size and defense intensity according to resource abundance, population density, season, and reproductive status. (Davies et al., 2012).

Human disturbance can alter territorial behavior by changing resource distribution, introducing artificial light and noise, or increasing encounters with people and domestic animals. Behavioral monitoring can therefore help identify habitat degradation and guide protected-area and corridor planning.

**8. Social Cooperation and Group Living:** Group living or social life has evolved in many taxa during evolution, because social interaction can provide important benefits that cannot achieved through individual living. Groups may help to improve predator detection, facilitate cooperative hunting, protect young ones from the predators, defend resources for better survival, or increase access to information. Examples shown by different wild animals where cooperative hunting is the key feature of carnivores, coordinated defense against any intruders in social mammals, and collective vigilance in birds and ungulates against predators. (Wilson, 1975; Silk, 2007).

Social cooperation in different species can also involve division of labor. In skilled social insects, individuals may show different characters as queen and king in honey bee specialize in reproduction, defensive activity shown by soldiers, brood care through nursing workers, or food collection. Among vertebrates, roles are generally more flexible and can depend on age, sex, dominance, kinship, and experience. Social learning allows individuals to acquire information from group members. (Wilson, 1975). Although group living is not universally beneficial in every animal. Competition for food and mates, disease transmission will help easily dispersion in group animals, social conflict is a very common phenomenon in same species, and increased visibility to predators can impose costs. The balance between benefits and costs depends strongly on ecological conditions (Silk, 2007).

**9. Communication and Information Transfer:** Communication in between wild animals allow to transfer information about identity, location or habitat, reproductive condition, territory ownership, danger, and resources. Signals may be visual, acoustic, chemical, tactile, or multimodal. Communication is central to social organization and can strongly influence survival and reproduction. (Bradbury & Vehrencamp, 2011).

Birdsong, mammalian voice calls, pheromones secreted by insects, scent marking in carnivores through the urination, and visual identification in reptiles are examples of specialized communication systems. Some signals in certain animals operate over long distances, whereas others function only at close range. Signal reliability depends on the environmental condition, receiver abilities, and potential costs bear by call creator (Bradbury & Vehrencamp, 2011). Communication also supports social learning. Animals can observe the behavior of others and modify their own actions. Artificial noise, light pollution, habitat fragmentation, and changes in group structure can interfere with normal signaling and social behavior. (Danchin et al., 2004).

**10. Foraging and Feeding Strategies:** Food acquisition is a fundamental determinant of survival. Wild animals use different foraging activities depending on population of prey distribution, food quality, competition, predation risk, and energy requirements. Some species specialize on particular foods, whereas others are generalists capable of exploiting many resources. (Stephens & Krebs, 1986).

Optimal foraging theory explains that different animals tend to choose certain behaviors that provide more favorable energetic returns in comparison to costs and risks faces by individuals. In general, decisions of the animals are influenced by incomplete information, learning, social cues, memory, and unpredictable environmental change. A predator may leave a profitable feeding site enriched with prey population if danger increases in specific area, while in case of herbivore may change its diet when preferred plants become scarce (Stephens & Krebs, 1986). Behavioral flexibility in feeding is increasingly important in human-modified ecosystems. Some species exploit agricultural fields, urban waste, artificial water sources, or supplementary feeding. Such flexibility may help populations persist, but it can also increase human-wildlife conflict and disease transmission. (Danchin et al., 2004).

**11. Reproductive Behaviour and Parental Care:** Reproductive behavior in wild animals includes selection of suitable mates, courtship behaviour shown by different animals, territorial displays, mating systems, nest construction for proper care, egg laying at suitable selected site, birth, and parental care of the young ones. These behaviors directly influence the reproductive success of the individual and are often closely linked with seasonal environmental conditions (Clutton-Brock, 1991). Courtship signals generated by specific species can communicate species identity, quality, readiness to reproduce, and competitive ability. After reproduction, parental care may include incubation, protection, feeding, grooming, carrying, teaching, and defense of offspring (Clutton-Brock, 1991).

Parental investment for future generation can substantially increase juvenile survival, but it also carries energetic and predation risks for adults. In certain species, cooperative breeding allows individuals other than the parents to assist in raising young. Changes in feeding habit and food availability, temperature, habitat structure, and human disturbance can disrupt breeding behavior. (Clutton-Brock, 1991).

**12. Behavioral Responses to Environmental Change:** Animals increasingly affected with environmental changes and variation caused by climate variability, habitat fragmentation due to anthropogenic activity, pollution, invasive species, artificial light, traffic, noise, and direct human presence. Behavior can provide a rapid mechanism for responding to these external pressures. Animals may shift their activity times, alter movement routes, change diet, modify habitat use, or become more tolerant of human presence (Tuomainen & Candolin, 2011). Behavioral plasticity can buffer populations against short-term environmental change, but plasticity has limits. If environmental conditions change faster than animals can respond, or if alternative habitats are unavailable, behavioral adjustment may not prevent population decline (Tuomainen & Candolin, 2011).

Climate change may alter the timing and season of migration, breeding in animals, emergence from dormancy condition, and food availability. If interacting species respond at different rates with changing environmental conditions, ecological mismatches can arise. Monitoring behavior alongside environmental variables can help detect such mismatches early (Tuomainen & Candolin, 2011).

**13. Importance for Wildlife Conservation:** Behavioral habit and information are valuable in wildlife conservation because population counts alone may not reveal real data changes in ecological condition. Alterations in vigilance and observational activity, movement of the individuals, feeding, breeding, vocal activity, habitat selection, or social structure can provide early indications of disturbance (Berger-Tal et al., 2011). Conservation of the wild animals and planning to save them can incorporate behavior by protecting migration corridors, breeding territories, nesting sites, roosting locations, and seasonal feeding grounds. Reducing wild animals' disturbance during their sensitive periods can be as important as protecting physical habitat. Wildlife crossings, quiet zones, lighting controls, and restoration of natural food and water resources can support normal behavioral patterns. (Berger-Tal et al., 2011).

Behavioral studies also help for upcoming generations for proper manage and help in less human-wildlife conflict. After understanding when and why animals enter agricultural fields or urban areas can guide non-lethal interventions. Effective conservation should aim not only to keep animals alive but to maintain ecological opportunities for feeding, reproduction, movement, and social organization. (Berger-Tal et al., 2011).

**14. Limitations and Research Gaps:** Behavior is difficult to measure consistently because it varies with individual personality, age, sex, season, social context, and immediate environmental conditions. Short observation periods may

therefore produce misleading conclusions for conservation of wild animals. Researchers should use standardized protocols, adequate sample sizes for study, repeatable observations, and appropriate statistical tools for proper data uses (Martin & Bateson, 2007).

Utilization of technological approaches such as camera traps, GPS telemetry, acoustic monitoring, drones, automated image recognition, and biologging can expand the scale of behavioral research. However, these tools utilization require careful validation of technology and may introduce observer or device effects. Ethical considerations are important when instruments are attached to animals or observations occur near breeding sites (Martin & Bateson, 2007; Berger-Tal et al., 2011). Future research related to behavioral studies should integrate behavioral ecology with physiology, genetics, landscape ecology, and conservation biology. Long-term studies are needed to determine whether behavioral flexibility improves population persistence under climate change and habitat fragmentation (Berger-Tal et al., 2011).

**15. Conclusion:** Behavioral adaptations in animals are central to the survival of wild animals. Migration of the animals, dormancy, camouflage character, antipredator behavior, territoriality, social cooperation, communication, foraging, and parental care allow animals to respond to complex ecological challenges. These strategies are shaped by natural selection but are also influenced by learning, experience, social relationships, and environmental conditions (Alcock, 2013; Davies et al., 2012). Every behavior of animals involves costs, benefits, constraints, and trade-offs. Animals capable of adjusting behavior to changing conditions may have important advantages under environmental variability. (Krebs & Davies, 1993). For wildlife conservation in natural environment, behavior should therefore be treated as an essential component of ecological assessment. Protecting habitats from the anthropogenic impact, movement routes, breeding areas, and social environments can help maintain the behaviors that sustain populations. Continued research into animal behavior will improve understanding of ecosystem functioning and support evidence-based wildlife management. (Berger-Tal et al., 2011).

## References:

1. Alcock, J. (2013). *Animal Behavior: An Evolutionary Approach*. Sinauer Associates.
2. Begon, M., Townsend, C. R., & Harper, J. L. (2006). *Ecology: From Individuals to Ecosystems*. Blackwell Publishing.
3. Campbell, N. A. et al. (2018). *Campbell Biology*. Pearson.
4. Dawkins, M. S. (1995). *Unravelling Animal Behaviour*. Longman.
5. Krebs, J. R., & Davies, N. B. (1993). *An Introduction to Behavioural Ecology*. Blackwell Scientific Publications.
6. Odum, E. P., & Barrett, G. W. (2005). *Fundamentals of Ecology*. Thomson Brooks/Cole.
7. Smith, T. M., & Smith, R. L. (2015). *Elements of Ecology*. Pearson.
8. Wilson, E. O. (1975). *Sociobiology: The New Synthesis*. Harvard University Press.
9. Clutton-Brock, T. H. (1991). *The Evolution of Parental Care*. Princeton University Press.
10. Fryxell, J. M., Sinclair, A. R. E., & Caughley, G. (2014). *Wildlife Ecology, Conservation, and Management*. Wiley-Blackwell.
11. Primack, R. B. (2014). *Essentials of Conservation Biology*. Sinauer Associates.
12. Ricklefs, R. E., & Relyea, R. (2014). *The Economy of Nature*. W. H. Freeman.
13. National Research Council. (2003). *Using Science to Improve the Nation's Wildlife Management*. National Academies Press.
14. Bolhuis, J. J., & Giraldeau, L.-A. (Eds.). (2005). *The Behavior of Animals: Mechanisms, Function, and Evolution*. Blackwell.
15. Danchin, E., Giraldeau, L.-A., & Cézilly, F. (2004). *Behavioural Ecology: An Evolutionary Perspective on Behaviour*. Oxford University Press.
16. Davies, N. B., Krebs, J. R., & West, S. A. (2012). *An Introduction to Behavioural Ecology* (4th ed.). Wiley-Blackwell.
17. Dingle, H., & Drake, V. A. (2007). What is migration? *BioScience*, 57(2), 113–121.

18. Berthold, P. (2001). *Bird Migration: A General Survey* (2nd ed.). Oxford University Press.
19. Geiser, F. (2013). Hibernation. *Current Biology*, 23(7), R188–R193.
20. Ruf, T., & Geiser, F. (2015). Daily torpor and hibernation in birds and mammals. *Biological Reviews*, 90(2), 891–926.
21. Ruxton, G. D., Sherratt, T. N., & Speed, M. P. (2004). *Avoiding Attack*. Oxford University Press.
22. Lima, S. L., & Dill, L. M. (1990). Behavioral decisions made under the risk of predation. *Canadian Journal of Zoology*, 68(4), 619–640.
23. Caro, T. (2005). *Antipredator Defenses in Birds and Mammals*. University of Chicago Press.
24. Silk, J. B. (2007). The adaptive value of sociality in mammalian groups. *Philosophical Transactions of the Royal Society B*, 362, 539–559.
25. Bradbury, J. W., & Vehrencamp, S. L. (2011). *Principles of Animal Communication* (2nd ed.). Sinauer Associates.
26. Stephens, D. W., & Krebs, J. R. (1986). *Foraging Theory*. Princeton University Press.
27. Tuomainen, U., & Candolin, U. (2011). Behavioural responses to human-induced environmental change. *Biological Reviews*, 86(3), 640–657.
28. Berger-Tal, O., et al. (2011). Integrating animal behavior and conservation biology: A conceptual framework. *Conservation Biology*, 25(1), 1–9.
29. Martin, P., & Bateson, P. (2007). *Measuring Behaviour* (3rd ed.). Cambridge University Press.
30. Mayr, E. (2004). *What Makes Biology Unique?* Cambridge University Press.
31. Wilcove, D. S., & Wikelski, M. (2008). Going, going, gone: Is animal migration disappearing? *PLoS Biology*, 6(7), e188.
32. Palmer MS, Packer C. Reactive anti-predator behavioral strategy shaped by predator characteristics. *PLoS One*. 2021 Aug 18;16(8): e0256147. doi: 10.1371/journal.pone.0256147. PMID: 34407141; PMCID: PMC8372962.
33. Caro TM. Ungulate antipredator behaviour: preliminary and comparative data from African bovids. *Behaviour*. 1994;189–228.
34. LaGory KE. The influence of habitat and group characteristics on the alarm and flight response of white-tailed deer. *Anim. Behav.* 1987;35(1): 20–25.
35. Hasson O. Pursuit-deterrent signals: communication between prey and predator. *Trends. Ecol. Evol.* 1991;6(10): 325–329. doi: 10.1016/0169-5347(91)90040-5
36. Jarman P. The social organisation of antelope in relation to their ecology. *Behaviour* 1974; 48(1–4): 215–267.
37. Michael A Fedak Ben Wilson and Patrick Pomeroy; *Reproductive Behavior*; December 2009; DOI: 10.1016/B978-0-12-373553-9.00217-0; book: *Encyclopedia of Marine Mammals*
38. Kovacks K; 2009; Maternal behaviour and early behavioural ontogeny of grey seals (*Halichoerus grypus*) on the Isle of May, UK; *Journal of Zoology* 213(4):697 – 715; DOI: 10.1111/j.1469-7998.1987.tb03735.x
39. Anholt RRH, O'Grady P, Wolfner MF, Harbison ST. Evolution of Reproductive Behavior. *Genetics*. 2020 Jan;214(1):49-73. doi: 10.1534/genetics.119.302263. PMID: 31907301; PMCID: PMC6944409.
40. Alexandra Alvergne, Vedrana Höggqvist Tabor; Is Female Health Cyclical? Evolutionary Perspectives on Menstruation; *Trends in Ecology & Evolution*; Volume 33, Issue 6, 2018, Pages 399-414, ISSN 0169-5347, <https://doi.org/10.1016/j.tree.2018.03.006>.
41. Lima SL, Dill LM. Behavioral decisions made under the risk of predation: a review and prospectus. *Can. J. Zool.* 1990;68(4): 619–640.
42. J.O. Wolff and J.A. Peterson 2010; An offspring-defense hypothesis for territoriality in female mammals; May 2010; *Ethology Ecology & Evolution* 10(3):227-239; DOI: 10.1080/08927014.1998.9522854
43. García-Ruiz I, Rubenstein DR. Fitness drivers of division of labor in vertebrates. *Elife*. 2026 Apr 8;14:RP105501. doi: 10.7554/eLife.105501. PMID: 41948805; PMCID: PMC13061414.



44. Lynch RF. Parents face quantity-quality trade-offs between reproduction and investment in offspring in Iceland. *R Soc Open Sci.* 2016 May 18;3(5):160087. doi: 10.1098/rsos.160087. PMID: 27293787; PMCID: PMC4892449.
45. Santos ESA, Nakagawa S. 2012. The costs of parental care: a meta-analysis of the trade-off between parental effort and survival in birds. *J. Evol. Biol.* 25, 1911–1917. (doi:10.1111/j.1420-9101.2012. 02569.x
46. Helm B, Liedvogel M. Avian migration clocks in a changing world. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol.* 2024 Jul;210(4):691-716. doi: 10.1007/s00359-023-01688-w. Epub 2024 Feb 2. PMID: 38305877; PMCID: PMC11226503.
47. Able KP, Belthoff JR. Rapid ‘evolution’ of migratory behaviour in the introduced house finch of eastern North America. *Proc Biol Sci.* 1998;265(1410):2063–2071. doi: 10.1098/rspb.1998.0541.
48. Åkesson S, Helm B. Endogenous programs and flexibility in bird migration. *Front Ecol Evol.* 2020;8(78):1–20. doi: 10.3389/fevo.2020.00078.
49. Åkesson S, Ilieva M, Karagicheva J, Rakhimberdiev E, Tomotani B, Helm B. Timing avian long-distance migration: from internal clock mechanisms to global flights. *Philos Trans R Soc Lond B Biol Sci.* 2017; 372:1734. doi: 10.1098/rstb.2016.0252.
50. Appenroth D, Wagner GC, Hazlerigg DG, West AC. Evidence for circadian-based photoperiodic timekeeping in Svalbard ptarmigan, the northernmost resident bird. *Curr Biol.* 2021 doi: 10.1016/j.cub.2021.04.009.
51. Archer LC, Hutton SA, Harman L, O'Grady MN, Kerry JP, Poole WR, Gargan P, McGinnity P and Reed TE (2019) The Interplay Between Extrinsic and Intrinsic Factors in Determining Migration Decisions in Brown Trout (*Salmo trutta*): An Experimental Study. *Front. Ecol. Evol.* 7:222. doi: 10.3389/fevo.2019.00222.